



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/725, 38/00, 31/445, 39/395, 39/40, 39/42	A1	(11) International Publication Number: WO 98/23279 (43) International Publication Date: 4 June 1998 (04.06.98)
(21) International Application Number: PCT/US97/21440 (22) International Filing Date: 24 November 1997 (24.11.97) (30) Priority Data: 60/031,878 27 November 1996 (27.11.96) US (71) Applicant (for all designated States except US): RHONE-POULENC RORER PHARMACEUTICALS INC. [US/US]; Legal Patents, Mail Stop 3C43, P.O. Box 5093, Collegeville, PA 19426-0997 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): UZAN, Andre [FR/FR]; 35, avenue Victor Hugo, F-75116 Paris (FR). CU-RAUDEAU, Alain, H. [FR/US]; 501 Brighton Circle, Devon, PA 19333 (US). LEADLEY, Robert, J. [US/US]; 6007 Indian Woods Lane, Collegeville, PA 19426 (US). DUNWIDDIE, Christopher, T. [US/US]; 3779 Trewithen Lane, Carmel, IN 46032 (US). PERRONE, Mark, H. [US/US]; 406 Hampton Court, Chalfont, PA 18914 (US). (74) Agents: PARKER, Raymond, S. et al.; Rhone-Poulenc Rorer Pharmaceuticals Inc., Mail Stop 3C43, P.O. Box 5093, Collegeville, PA 19426-0997 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PHARMACEUTICAL COMPOSITION COMPRISING A COMPOUND HAVING ANTI-Xa ACTIVITY AND A PLATELET AGGREGATION ANTAGONIST COMPOUND		
(57) Abstract <p>The invention is directed to pharmaceutical composition comprising a compound having anti-Xa activity, a platelet aggregation antagonist compound and a pharmaceutically acceptable carrier. The invention is also directed to a method of treating or preventing a thrombogenic condition associated with a thrombosis related ischemic disorder in a patient comprising administering to said patient pharmaceutically effective amounts of a compound having anti-Xa activity and a platelet aggregation antagonist compound. In addition, this invention is directed to the use of pharmaceutically effective amounts of a compound having anti-Xa activity and a platelet aggregation antagonist compound in the preparation of a medicament for treating or preventing a physiological condition associated with thrombosis related ischemic disorder. Furthermore, this invention is directed to a kit for treating or preventing a physiological condition associated with thrombosis related ischemic disorder, said kit comprising a plurality of separate containers, wherein at least one of said containers contains a compound having anti-Xa activity and at least another of said containers contains a platelet aggregation antagonist compound, and said containers optionally contain a pharmaceutical carrier.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PHARMACEUTICAL COMPOSITION COMPRISING A COMPOUND
HAVING ANTI-Xa ACTIVITY AND A PLATELET AGGREGATION ANTAGONIST
COMPOUND

5 Field of the Invention

This invention is directed to a pharmaceutical composition comprising an anti-Xa activity compound and a platelet aggregation antagonist compound which exhibits unexpectedly efficacious activity for treating or preventing a physiological condition associated with a thrombosis related ischemic disorder in a patient. The invention is also directed to a method of treating or preventing a thrombogenic condition associated with a thrombosis related ischemic disorder in a patient comprising administering pharmaceutically effective amounts of an anti-Xa activity compound and a platelet aggregation antagonist compound.

15 Platelet fibrinogen receptor antagonists have been shown to be effective agents for inhibiting platelet-dependent thrombus formation in animal models of coronary thrombosis. Likewise, results from clinical studies have shown that platelet fibrinogen receptor antagonists reduce the composite incidence of major ischemic events when administered to high-risk patients undergoing percutaneous transluminal coronary angioplasty. However, the therapeutic window is very narrow for this class of compounds, in part, because the high degree of inhibition of ex vivo platelet aggregation required for antithrombotic efficacy is often associated with a marked increase in template bleeding time, which is a marker of an undesirable bleeding complications.

25 Low molecular weight heparins (LMWHs) and heparinoid compounds (HCs) have been used effectively over the past few years for the prevention and treatment of venous thrombosis and the associated thromboembolism. However, LMWHs and HCs are gradually pervading into the treatment repertoire for arterial thrombotic indications. Preliminary results favoring the use of LMWH or HCs over unfractionated heparin in arterial thrombotic indications are supported by several pharmacodynamic, pharmacokinetic, and mechanistic differences between these two classes of compounds. For example, reliable and safe anticoagulation can be achieved with LMWHs or HCs by subcutaneous dosing without monitoring. Compared to heparin, LMWHs and HCs have a higher bioavailability, a relatively long half-life, and appear to have a safer profile. In addition, LMWHs are more resistant than heparin to neutralization by platelet factor 4, which is released from activated platelets, presumably at the site of arterial thrombosis.

35 Haemostasis, the biochemistry of blood coagulation, is an extremely complex phenomena whereby normal whole blood and body tissue spontaneously arrest bleeding from injured blood vessels. Effective haemostasis requires the combined activity of vascular, platelet and plasma factors

as well as a controlling mechanism to prevent excessive clotting. Defects, deficiencies, or excesses of any of these components can lead to hemorrhagic or thrombotic consequences.

Platelet adhesion, spreading and aggregation on extracellular matrices are central events in thrombus formation. These events are mediated by a family of adhesive glycoproteins, i.e., fibrinogen, fibronectin, and von Willebrand factor. Fibrinogen is a co-factor for platelet aggregation, while fibronectin supports platelet attachments and spreading reactions, and von Willebrand factor is important in platelet attachment to and spreading on subendothelial matrices. The binding sites for fibrinogen, fibronectin and von Willebrand factor have been located on the platelet membrane protein complex known as glycoprotein IIb/IIIa.

Adhesive glycoproteins, like fibrinogen, do not bind with normal resting platelets. However, when a platelet is activated with an agonist such as thrombin or adenosine diphosphate, the platelet changes its shape, perhaps making the GPIIb/IIIa binding site accessible to fibrinogen. Blocking the fibrinogen receptor, thus inhibits platelet aggregation and subsequent thrombus formation, and is useful for the prevention and treatment of pathological thrombogenic conditions, such as stroke, peripheral arterial occlusive disease, disseminated intravascular coagulation, and acute coronary syndromes such as unstable angina and myocardial infarction.

SUMMARY OF THE INVENTION

The invention is also directed to pharmaceutical composition comprising a compound having anti-Xa activity, a platelet aggregation antagonist compound and a pharmaceutically acceptable carrier. The invention is also directed to a method of treating or preventing a thrombogenic condition associated with a thrombosis related ischemic disorder in a patient comprising administering to said patient pharmaceutically effective amounts of a compound having anti-Xa activity and a platelet aggregation antagonist compound.

BRIEF DESCRIPTION OF THE FIGURES

FIGURE 1 represents information regarding assorted blood sampling /hemodynamic measurements during administration of different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, N-(n-butylsulfonyl)-4-(piperidin-4-ylbutyloxy)-L-phenylalanine (BSPBPA), and composition thereof over time.

FIGURE 2 represents a graph of the number of cyclic flow reductions (CFRs) for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, BSPBPA, and composition thereof over time.

FIGURE 3 represents a graph of the activated partial thromboplastin time (APTT) for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, BSPBPA, and composition thereof over time.

5 FIGURE 4 represents a graph of the prothrombin time (PT) for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, BSPBPA, and composition thereof over time.

10 FIGURE 5 represents a graph of the Anti-Xa activity for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, BSPBPA, and composition thereof over time.

15 FIGURE 6 represents a graph of the Anti-IIa activity for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, BSPBPA, and composition thereof over time.

20 FIGURE 7 represents a graph of the template bleeding time for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, BSPBPA, and composition thereof over time.

 FIGURE 8 represents a graph of the platelet count for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, BSPBPA, and composition thereof over time.

25 FIGURE 9 represents a graph of collagen-induced ex-vivo platelet aggregation for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, BSPBPA, and composition thereof over time.

30 FIGURE 10 represents a graph of ADP-induced ex-vivo platelet aggregation for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, BSPBPA, and composition thereof over time.

35 FIGURE 11 represents a graph of arachidonic acid-induced ex-vivo platelet aggregation for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, BSPBPA, and composition thereof over time.

 FIGURE 12 represents a graph of thrombin-induced ex-vivo platelet aggregation for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, BSPBPA, and composition thereof over time.

FIGURE 13 represents information regarding assorted blood sampling /hemodynamic measurements during administration of different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, N-[N-[N-(4-(piperidin-4-yl)butanoyl)-N-ethylglycyl]aspartyl]-L- β -cyclohexyl alanine amide (PBGACA), and composition thereof over time.

FIGURE 14 represents a graph of the number of cyclic flow reductions (CFRs) for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, PBGACA, and composition thereof over time.

FIGURE 15 represents a graph of the activated partial thromboplastin time (APTT) for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, PBGACA, and composition thereof over time.

FIGURE 16 represents a graph of the prothrombin time (PT) for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, PBGACA, and composition thereof over time.

FIGURE 17 represents a graph of the Anti-Xa activity for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, PBGACA, and composition thereof over time.

FIGURE 18 represents a graph of the Anti-IIa activity for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, PBGACA, and composition thereof over time.

FIGURE 19 represents a graph of the template bleeding time for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, PBGACA, and composition thereof over time.

FIGURE 20 represents a graph of the platelet count for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, PBGACA, and composition thereof over time.

FIGURE 21 represents a graph of collagen-induced ex-vivo platelet aggregation for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, PBGACA, and composition thereof over time.

FIGURE 22 represents a graph of ADP-induced ex-vivo platelet aggregation for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, PBGACA, and composition thereof over time.

FIGURE 23 represents a graph of arachidonic acid-induced ex-vivo platelet aggregation for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, PBGACA, and composition thereof over time.

FIGURE 24 represents a graph of thrombin-induced ex-vivo platelet aggregation for different concentrations of the anti-Xa activity compound, Enoxaparin, the platelet aggregation antagonist compound, PBGACA, and composition thereof over time.

DETAILED DESCRIPTION OF THE INVENTION

As used above, and throughout the description of the invention, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

Definitions

"Patient" includes both human and other mammals.

"Effective amount" is meant to describe an amount of composition according to the present invention effective in producing the desired therapeutic effect.

"Platelet aggregation antagonist compound" (PAAC) means a compound that binds to the platelet GPIIb/IIIa receptor (GPIIb/IIIa receptor antagonist) and competitively inhibits binding of fibrinogen, fibronectin and von Willebrand factor as well as inhibits aggregation of activated platelets.

"Anti-Xa activity compound" means a heparinoid compound or low molecular weight heparin (LMWH), or synthetic derivatives thereof.

Preferred Embodiments

According to a preferred embodiment of the invention, the following references, incorporated herein by reference, describes useful PAACs: Lynch et al. J. Pharm. Expt. Thera. 272(1) 20 (1995); Kereiakes et al. JACC 27(3), 536 (1996); Peerlinck et al. Circulation 88(4), 1512 (1993); Barrett et al. Clin. Pharmacol. Ther. 56(4) 377 (1994); Cook et al. Thromb. Haemostas. 70(5), 838 (1993); Plow, et al., Proc. Natl. Acad. Sci. USA 82, 8057-61 (1985); Ruggeri, et al., Proc. Natl. Acad. Sci. USA 5708-12 (1986); Ginsberg, et al., J. Biol. Chem. 260, 3931-36 (1985); and Gartner, et al., J. Biol. Chem. 260, 11,891-94 (1987); Plow, E.F., et al., Proc. Natl. Acad. Sci. USA 79, 3711-3715 (1982); Tjoeng, et al., U.S. Patent Nos. 5,037,808, 4,879,313 and 4,992,463; Adams, et al., U.S. Patent No. 4,857,508; Haverstick, D.M., et al., Blood 66(4), 946-952 (1985); Topol et al., The Lancet, 343, 881 (1994), French Application No. 86/17507; Zimmerman, et al., U.S. Patent No. 4,683,291; European Application Publication No. 0 319 506; U.S. Patent No. 5,023,233; U.S. Patent

No. 4,952,562; International Publication No. WO 91/04746; United States Application Serial No. 5,086,069; International Publication No. WO 92/13117; U.S. Pat. No. 5,053,392; U.S. Pat. No. 5,064,814; U.S. Pat. No. 5,051,405; European Patent Application 0479,481; European Patent Application 0478,362; U.S. Pat. No. 5,292,756; International Publication No. WO 95/10295; and
5 International Publication No. WO 89/11538. More preferred PAACs are those disclosed in International Publication No. WO 89/11538, International Publication No. WO 95/10295 or U.S. Pat. No. 5,292,756; further preferred are Reopro[®] (abciximab), N-[N-[N-(4-(piperidin-4-yl)butanoyl)-N-ethylglycyl]aspartyl]-L- β -cyclohexyl alanine, N-[N-[N-(4-(piperidin-4-yl)butanoyl)-N-ethylglycyl]aspartyl]-L- β -cyclohexyl alanine amide (PBGACA) or N-(n-butylsulfonyl)-4-(piperidin-
10 4-ylbutyloxy)-L-phenylalanine (BSPBPA).

According to a preferred embodiment of the invention, the following references, incorporated herein by reference, describes useful LMWHs according to the invention and methods for preparing the LMWHs: Eur. Pat. No. 0014184; Medicinal Research Reviews 12(4), 373 (1992);
15 Drugs of the Future 12(1), 45 (1987); International Publication No. WO 92/19249; U.S. Pat. No. 4,692,435; Barrowcliffe, Thromb. Res. 12, 27-36 (1977); Eur. Pat. Appln. No. 37319; Eur. Pat. Appln. No. 76279; U.S. Pat. No. 4,804,652; WO81/3276; Eur. Pat. Appln. No. 244235, Eur. Pat. Appln. No. 244236; U.S. Pat. No. 4,486,420; U.S. Pat. No. 4,692,435; U.S. Pat. No. 4,826,827; U.S. Pat. No. 3,766,167; Eur. Pat. Appln. No. 40144; Eur. Pat. Appln. No. 347588, Eur. Pat. Appln. No.
20 380943; U.S. Pat. No. 4533549; U.S. Pat. No. 4,629,699; Eur. Pat. Appln. No. 269981. For example, an anti-Xa activity compound may be produced as follows: enrichment by fractionation by ethanol and/or molecular sieving, e.g., gel filtration or membrane filtration of the LMWH present in standard heparin and controlled chemical (by nitrous acid, β -elimination or periodate oxidation) or enzymatic (by heparinases) depolymerization. The conditions for depolymerization can be carefully
25 controlled to yield products of desired molecular weights. Nitrous acid depolymerization is commonly used. Also employed is depolymerization of the benzylic ester of heparin by β -elimination, which yields the same type of fragments as enzymatic depolymerization using heparinases. LMWH with low anticoagulant activity and retaining basic chemical structure are prepared by depolymerization using periodate oxidation or by removing the antithrombin-binding
30 fraction of LMWH, or prepared by other methods, using immobilized antithrombin for adsorption.

Further preferred is a LMWH having an average molecular weight of about 3000 to about 6500. Commercially available LMWHs useful according to the invention include the following: Clexane[®]/Klexane[®]/Lovenox[®] (Enoxaparin (ENOX)) having an average molecular mass of $4500 \pm$
35 1000 Dalton (Da), molecular mass distribution comprising components of <2000 Da ($16.0 \pm 4.0\%$) and 2000 to 8000 Da ($78.0 \pm 10.0\%$), Anti-Xa activity (IU/mg on dry basis) of 90 to 125 , and an Anti-Xa/Anti-IIa ratio of 3.3 to 5.3 ; Fraxiparin[®] (Nardroparin) having an average molecular mass of 4300 ± 700 Da, molecular mass distribution comprising components of <2000 Da ($<15\%$), 2000 to 4000 Da ($45 \pm 10\%$) and 2000 to 8000 Da ($85 \pm 10\%$), Anti-Xa activity (IU/mg on dry basis) of 95
40 to 130 , and an Anti-Xa/Anti-IIa ratio of 2.5 to 4.0 ; Fragmin[®] (Dalteparin) having an average molecular mass of 6000 ± 400 Da, molecular mass distribution comprising components of <3000 Da

(5.0 to 13.0%) and >8000 Da (15.0 to 25.0%), Anti-Xa activity (IU/mg on dry basis) of 110 to 210, and an Anti-Xa/Anti-IIa ratio of 1.9 to 3.2; Embolex®/Monoembolex® (Certroparin) having an average molecular mass of 5200 ± 1000 Da, molecular mass distribution comprising components of <2000 Da (10 to 25%) and <8000 Da (75 to 90%), Anti-Xa activity (IU/mg on dry basis) of 80 to 120, and an Anti-Xa/Anti-IIa ratio of 1.5 to 2.5; Fluxum®/Minidaltan®/Lowhepa® (Parnaparin) having an average molecular mass of 5000 ± 1000 Da, molecular mass distribution comprising components of <3000 Da (20 to 30%) and 3000 to 8000 Da (50 to 60%), Anti-Xa activity (IU/mg on dry basis) of 75 to 110, and an Anti-Xa/Anti-IIa ratio of 2.0 to 3.0; Logiparin® (Tinzaparin) having an average molecular mass of 3400 to 5600 Da, molecular mass distribution comprising components of <2000 Da (2.0 to 16.0%), 2000 to 4000 Da ($66.0 \pm 6.0\%$) and >8000 Da (12.0 to 38.0%), Anti-Xa activity (IU/mg on dry basis) of >70, and an Anti-Xa/Anti-IIa ratio of 1.5 to 2.5; Clivarine® (Reviparin) and Normiflo® (ardeparin/RD heparin/RDH). A preferred LMWH according to the invention is prepared according to the procedures disclosed in U.S. Pat. No. 4,486,420 or U.S. Pat. No. 4,692,435; more preferred enoxaparin.

According to another preferred embodiment of the invention, an anti-Xa activity compound is a heparinoid compound. The following references, incorporated herein by reference, describe a useful heparinoid compound according to the invention and methods for preparing the heparinoid compound: U.S. Pat. No. 4,438,108, Eur. Pat. No. EP 66908, Zammit et al. *Thromb. Haemostas.* 71(6), 759 (1994), and Gent et al. *Circulation* 93, 80 (1996). A preferred commercially available heparinoid compound useful according to the invention is Orgaran® (Danaparoid) having an average molecular mass of about 6500 Da, Anti-Xa activity (IU/mg on dry basis) of about 10, and an Anti-Xa/Anti-IIa ratio of about 28.

Preferred thrombogenic conditions treatable or preventable according to the invention include stroke, atherosclerosis, angiogenesis, thrombosis, thromboembolic conditions such as deep venous thrombosis, pulmonary embolism or thrombophlebitis, disseminated intravascular coagulation or thromboembolic syndromes associated with cancer, sepsis or obstetrical complications, peripheral arterial occlusive disease, and acute coronary syndromes such as unstable angina and myocardial infarction, hemodialysis, or extra-corporeal circulation requirement associated with a surgical procedure or tissue damage caused by phospholipases A2 (PLA₂); more preferred is unstable angina and myocardial infarction.

Another preferred embodiment according to the invention useful in the course of a medical procedure wherein there is a potential for an occurrence pathological thrombogenic condition may occur, such as during coronary artery bypass surgery or percutaneous transluminal coronary angioplasty with or without placement of an intracoronary stent; more preferred is percutaneous transluminal coronary angioplasty with or without placement of an intracoronary stent.

Another preferred embodiment according to the invention is the use of pharmaceutically effective amounts of a compound having anti-Xa activity and a platelet aggregation antagonist

compound in the preparation of a medicament for treating or preventing a physiological condition associated with thrombosis related ischemic disorder.

In the treatment or prevention method according to the invention the anti-Xa activity compound and platelet aggregation antagonist compound may be administered in different ways, such as in combination therapies optionally employing medical procedures. For example, the anti-Xa activity compound and platelet aggregation antagonist compound may be administered to a patient concomitantly or at different times provided that they are administered such that at some period of time there are pharmaceutically effective amounts of both compounds present in the patient such that a therapeutic effect according to the invention results.

Thus, it is a further object of the invention to provide a kit for treating or preventing a physiological condition associated with thrombosis related ischemic disorder, said kit comprising a plurality of separate containers, wherein at least one of said containers contains a compound having anti-Xa activity and at least another of said containers contains a platelet aggregation antagonist compound, and said containers optionally contain a pharmaceutical carrier, which kit may be effectively utilized for carrying out combination therapies according to the invention. A further embodiment for a kit would be wherein of said containers at least one of said containers should contain the compound having anti-Xa activity without the presence of the platelet aggregation antagonist compound, and at least another of said containers should contain the platelet aggregation antagonist compound without the presence of the compound having anti-Xa activity.

In practice, the anti-Xa activity compound and platelet aggregation antagonist compound may be administered parenterally, topically, rectally, transdermally, intrapulmonary or orally, but they are preferably administered parenterally and/or orally.

Suitable compositions containing the compounds used according to the invention may be prepared by conventional means. For example, the compounds used according to the invention may be dissolved or suspended in a suitable carrier.

The compounds used according to the invention should be presented in forms permitting administration by the most suitable route, and the invention also relates to a pharmaceutical composition containing the compounds used according to the invention which are suitable for use in human or veterinary medicine. These compositions may be prepared according to the customary methods, using one or more pharmaceutically acceptable carrier, which comprise adjuvants or excipients. The adjuvants comprise, inter alia, diluents, sterile aqueous media and the various non-toxic organic solvents. The compositions may be presented in the form of tablets, pills, capsules, lozenges, troches, hard candies, granules, powders, aqueous solutions or suspensions, injectable solutions, elixirs or syrups, powders, solution or suspension for intrapulmonary administration and

can contain one or more agents chosen from the group comprising sweeteners, flavorings, colorings, or stabilizers in order to obtain pharmaceutically acceptable preparations.

The choice of vehicle and the content of compounds used according to the invention in the vehicle are generally determined in accordance with the solubility and chemical properties of the compounds, the particular mode of administration and the provisions to be observed in pharmaceutical practice. For example, excipients such as sterile water, Ringer's solution, lactose, sodium citrate, isotonic saline solutions (monosodium or disodium phosphate, sodium, potassium, calcium or magnesium chloride, or mixtures of such salts), calcium carbonate and disintegrating agents such as starch, alginic acids and certain complex silicates combined with lubricants such as magnesium stearate, sodium lauryl sulfate and talc may be used for preparing tablets. To prepare a capsule, it is advantageous to use lactose and high molecular weight polyethylene glycols. When aqueous suspensions are used they can contain emulsifying agents or agents which facilitate suspension. Diluents such as sucrose, ethanol, polyethylene glycol, propylene glycol, glycerol and chloroform or mixtures thereof may also be used.

For parenteral administration, emulsions, suspensions or solutions of the compounds used according to the invention in vegetable oil, for example sesame oil, groundnut oil or olive oil, or aqueous-organic solutions such as water and propylene glycol, injectable organic esters such as ethyl oleate, as well as sterile aqueous solutions of the pharmaceutically acceptable salts, are useful. The solutions of the salts of the compounds used according to the invention are especially useful for administration by intramuscular, intravenous, intraarterial or subcutaneous injection or infusion techniques. The aqueous solutions, also comprising solutions of the salts in pure distilled water, may be used for intravenous administration with the proviso that their pH is suitably adjusted, that they are judiciously buffered and rendered isotonic with a sufficient quantity of glucose or sodium chloride and that they are sterilized by heating, irradiation or microfiltration.

The anti-Xa activity compound and platelet aggregation antagonist compound according to the invention may also be formulated in a manner which resists rapid clearance from the vascular (arterial or venous) wall by convection and/or diffusion, thereby increasing the residence time of the composition at the desired site of action. Depot useful according to the invention may be in a copolymer matrix, such as ethylene-vinyl acetate, or a polyvinyl alcohol gel surrounded by a Silastic shell. Alternatively, the anti-Xa activity compound and platelet aggregation antagonist compound may be delivered locally from a silicone polymer implanted in the adventitia.

An alternative approach for minimizing washout of the anti-Xa activity compound and platelet aggregation antagonist compound during percutaneous, transvascular delivery comprises the use of nondiffusible, drug-eluting microparticles. The microparticles may be comprised of a variety of synthetic polymers, such as polylactide for example, or natural substances, including proteins or

polysaccharides. Such microparticles enable strategic manipulation of variables including total dose of a drug and kinetics of its release. Microparticles can be injected efficiently into the arterial or venous wall through a porous balloon catheter or a balloon over stent, and are retained in the vascular wall and the periadventitial tissue for at least about two weeks. Formulations and methodologies for local, intravascular site-specific delivery of therapeutic agents are discussed in Reissen et al. (J. Am. Coll. Cardiol. 1994; 23: 1234-1244), the entire contents of which are hereby incorporated by reference.

The medium for the anti-Xa activity compound and platelet aggregation antagonist compound can also be a hydrogel which is prepared from any biocompatible or non-cytotoxic (homo or hetero) polymer, such as a hydrophilic polyacrylic acid polymer that can act as a drug absorbing sponge. Such polymers have been described, for example, in application WO93/08845, the entire contents of which are hereby incorporated by reference. Certain of them, such as, in particular, those obtained from ethylene and/or propylene oxide are commercially available.

In addition, the anti-Xa activity compound and platelet aggregation antagonist compound may be administered directly to the blood vessel wall by means of an angioplasty balloon which is coated with a hydrophilic film (for example a hydrogel), or by means of any other catheter containing an infusion chamber for the compounds, which can thus be applied in a precise manner to the site to be treated.

The percentage of the anti-Xa activity compound and platelet aggregation antagonist compound used according to the invention may be varied. The compounds should constitute a proportion such that a suitable dosage shall be obtained. Obviously, several unit dosage forms may be administered. The dose employed will be determined by the physician, and depends upon the desired therapeutic effect, the route of administration and the duration of the treatment, and the condition of the patient. In each particular case, the doses will be determined in accordance with the factors distinctive to the subject to be treated, such as age, weight, general state of health and other characteristics which can influence the efficacy of the medicinal product.

In the adult, the dosages of the PAAC are generally from about 0.0001 to about 50, preferably about 0.0001 to about 5, mg/kg body weight per day by inhalation, from about 0.001 to about 100, preferably 0.01 to 70, more especially 0.05 to 10, mg/kg body weight per day by oral administration, and from about 0.0001 to about 10, preferably 0.001 to 1, mg/kg body weight per day by intravenous administration.

In the adult, the dosages of the anti-Xa activity compound according to the invention is particularly useful in doses of about 10 to about 25,000 international units of anti-Xa activity.

The anti-Xa activity compound and platelet aggregation antagonist compound used according to the invention may be administered as frequently as necessary in order to obtain the

desired therapeutic effect. The dosage regimen in carrying out the method of this invention is that which insures maximum therapeutic response until improvement is obtained and thereafter the minimum effective level which gives relief. Some patients may respond rapidly to a higher or lower dose and may find much weaker maintenance doses adequate. Both short- and long-term treatments regimens are contemplated for the invention. Treatments at the rate of about 1 to about 4 doses per day are also contemplated, in accordance with the physiological requirements of each particular patient, bearing in mind, of course, that in selecting the appropriate dosages in any specific case, consideration must be given to the patient's weight, general health, age, and other factors which may influence response to the drug. Thus, for other patients, it will be necessary to prescribe not more than one or two doses per day.

The compounds of the present invention may also be formulated for use in conjunction with other therapeutic agents such as agents or in connection with the application of therapeutic techniques to address pharmacological conditions which may be ameliorated through the application of a compound of formula I, such as in the following:

The compounds of the present invention may be used in the treatment of restenosis post angioplasty using any device such as balloon, ablation or laser techniques. The compounds of the present invention may be used in the treatment of restenosis following stent placement in the vasculature either as 1) primary treatment for vascular blockage, or 2) in the instance where angioplasty using any device fails to give a patent artery. The compounds of the present invention may be used either orally, by parenteral administration or the compound could be applied topically through the intervention of a specific device or as a properly formulated coating on a stent device.

The compounds of the present invention may be used in the treatment of restenosis in combination with any anticoagulant, antiplatelet, antithrombotic or profibrinolytic agent. Often patients are concurrently treated prior, during and after interventional procedures with agents of these classes either in order to safely perform the interventional procedure or to prevent deleterious effects of thrombus formation. Some examples of classes of agents known to be anticoagulant, antiplatelet, antithrombotic or profibrinolytic agents include any formulation of thrombin inhibitors or Factor VIIa inhibitors. Some examples of classes of agents known to be anticoagulant, antiplatelet, antithrombotic or profibrinolytic agents include any formulation of aspirin, direct thrombin inhibitors, direct Factor Xa inhibitors, or Factor VIIa inhibitors.

The compounds of the present invention may be used in combination with any antihypertensive agent or cholesterol or lipid regulating agent in the treatment of restenosis or atherosclerosis concurrently with the treatment of high blood pressure or atherosclerosis. Some examples of agents that are useful in the treatment of high blood pressure include compounds of the following classes; beta-blockers, ACE inhibitors, calcium channel antagonists and alpha-receptor antagonists. Some examples of agents that are useful in the treatment of elevated cholesterol levels or

disregulated lipid levels include compounds known to be HMGCoA reductase inhibitors, compounds of the fibrate class,

The anti-Xa activity compound and platelet aggregation antagonist compound used according to the invention exhibit marked pharmacological activities according to tests described in the literature which tests results are believed to correlate to pharmacological activity in humans and other mammals. The following pharmacological test results are typical characteristics of the anti-Xa activity compound and platelet aggregation antagonist compound used according to the invention.

The following pharmacological tests evaluate the activity of anti-Xa activity compound and platelet aggregation antagonist compound used according to the invention. Those tests include the hemodynamics measurements during administration of different concentrations of an anti-Xa activity compound, a platelet aggregation antagonist compound and composition thereof over time (Figures 1 and 13). More specifically measurements were taken of cyclic flow reduction (Figures 2 and 14), activated partial thromboplastin time (Figures 3 and 15), prothrombin time (Figures 4 and 16), Anti-Xa activity (Figures 5 and 17), Anti-IIa activity (Figures 6 and 18), template bleeding time (Figures 7 and 19), platelet count (Figures 8 and 20), collagen-induced ex-vivo platelet aggregation (Figures 9 and 21), ADP-induced ex-vivo platelet aggregation (Figures 10 and 22), arachidonic acid-induced ex-vivo platelet aggregation (Figures 11 and 23), and thrombin-induced ex-vivo platelet aggregation (Figures 12 and 24).

Compositions of the present invention exhibit marked activity in the foregoing tests and are considered useful in the prevention and treatment of thrombosis associated with certain disease states. Antithrombotic activity in the ex vivo canine platelet aggregation assay is predictive of such activity in humans (see, for example, Catalfamo, J. L., and Dodds, W. Jean, "Isolation of Platelets from Laboratory Animals", Methods Enzymol. 169, Part A, 27 (1989)).

Materials and Methods

All procedures in this study are performed in compliance with the Animal Welfare Act Regulations and with the Guide for the Care and Use of Laboratory Animals (DHEW Publication No. NIH 85-23, 1985).

The test protocol as follows is an experimental model of unstable angina.

Mongrel dogs of either sex (15-21 kg) are anesthetized with sodium pentobarbital (30 mg/kg, i.v., with supplements given as needed), intubated, and ventilated using a Harvard respirator (Harvard Apparatus, S. Natick, MA). A tri-lumen catheter (SAFEDWELLplus, Becton Dickinson, Sandy, UT) is placed in the right femoral vein for the administration of test agents and supplemental anesthesia.

The right femoral artery is cannulated for measurement of arterial blood pressure and for obtaining blood samples.

A left thoracotomy is performed at the 5th intercostal space and the heart is suspended in a pericardial cradle. The left circumflex coronary artery (LCX) is isolated and dissected for a distance of 2 cm, ligating side branches when necessary. An electromagnetic flow probe (Carolina Medical Electronics, 501D) is placed on the vessel to monitor coronary blood flow and a snare ligature is placed on the distal portion of the vessel to produce a temporary mechanical occlusion which is used to aid in adjusting the degree of stenosis and to aid in validating zero flow measurements.

Distal to the flow probe, a Lexan[®] occluder is positioned for the purpose of creating a critical stenosis, which is confirmed by abolishment of the hyperemic response to a 10 sec mechanical occlusion of the vessel. The endothelium and vascular smooth muscle cells are damaged by compressing the vessel with a vascular clamp. These conditions result in platelet adhesion and aggregation at the damaged area, thus producing a gradual decrease in coronary blood flow. When flow reaches zero, the occluder is moved back and forth over the damaged area to mechanically dislodge the platelet-rich thrombus, thus restoring blood flow. This repetitive pattern of decreasing blood flow that is restored by mechanical disruption of the platelet thrombus is referred to as cyclic flow reductions (CFRs). The antithrombotic effect of the test agents is quantitated by comparing the number of CFRs that occurred during a 20 min control period with the number of CFRs per 20 min for three consecutive 20 min periods after drug administration. A significant reduction in the number of CFRs is taken to represent an antithrombotic effect.

Experimental Protocol

a. Protocol using BSPBPA

Thirty dogs are assigned to one of six adjunctive treatment groups. The compounds are administered as an intravenous bolus only (for N-(n-butylsulfonyl)-4-(piperidin-4-ylbutyloxy)-L-phenylalanine (BSPBPA) or saline vehicle) or as an intravenous bolus plus a constant intravenous infusion (for heparin, enoxaparin (ENOX), or saline vehicle). The treatment groups are: I) BSPBPA (30 µg/kg), II) BSPBPA (300 µg/kg), III) ENOX (0.5 mg/kg + 5 µg/kg/min) IV) ENOX (0.5 mg/kg + 5 µg/kg/min) plus BSPBPA (30 µg/kg), V) heparin (60 U/kg + 0.7 U/kg/min), and VI) heparin (60 U/kg + 0.7 U/kg/min) plus BSPBPA (30 µg/kg). All compounds are diluted in saline and bolus injections are made using a volume of 5 mL and constant infusions are made using a volume of 22 mL.

a. Protocol using PBGACA

Thirty dogs are assigned to one of six adjunctive treatment groups. The compounds for these experiments involves the administration of the following agents as a bolus and a constant infusion (for N-[N-[N-(4-(piperidin-4-yl)butanoyl)-N-ethylglycyl]aspartyl]-L- β -cyclohexyl alanine amide (PBGACA) or saline vehicle) or as an intravenous bolus plus a constant intravenous infusion (for heparin, ENOX, or saline vehicle. The treatment groups were: I) PBGACA (10 μ g/kg+ 0.15 μ g/kg/min), II) PBGACA (30 μ g/kg + 0.15 μ g/kg/min), III) ENOX (0.5 mg/kg + 5 μ g/kg/min) IV) ENOX (0.5 mg/kg + 5 μ g/kg/min) plus PBGACA (10 μ g/kg+ 0.15 μ g/kg/min), V) heparin (60 U/kg + 0.7 U/kg/min), and VI) heparin (60 U/kg + PBGACA (10 μ g/kg+ 0.15 μ g/kg/min), . All compounds are diluted in saline and bolus injections are made using a volume of 5 mL and constant infusions were made using a volume of 22 mL.

After consistent CFRs are established for at least 20 minutes, compounds are administered as described above and blood flow is monitored during the one hour infusion period. Arterial blood samples are drawn before administration of test agents (control sample) and at 5, 10, 30, and 60 min after administration of compounds. Blood samples are drawn on 1/10 volume of 3.8% trisodium citrate and are used for ex vivo platelet aggregation, platelet count, anti-Xa levels, anti-IIa levels, and coagulation time assays (activated partial thromboplastin time, APTT, and prothrombin time, PT). Blood samples (4.5 mL) obtained for analysis of anti-Xa and anti-IIa levels are collected in chilled syringes containing 0.5 mL trisodium citrate and immediately placed on ice. Mean arterial blood pressure, heart rate and EKG are recorded for the duration of the protocol (Grass polygraph, Model 7D, Grass Instruments, MA).

Coagulation times and template bleeding time. Activated partial thromboplastin time (APTT) and prothrombin time (PT) are measured using a Microsample Coagulation Analyzer (MCA210, Bio Data Corp, Horsham, PA) and Dade[®] reagents (Thromboplastin-C Plus and Actin[®] FS Activated PTT reagent, Baxter Diagnostics, Inc., Deerfield, IL).

The APTT is the most widely used method for monitoring intravenous heparin anticoagulation therapy. It is also a fundamental screening test for deficiencies or abnormalities of the intrinsic coagulation factors: VIII, IX, XI, XII, and factors common to both the intrinsic and extrinsic pathways: I (fibrinogen), II, V, X. When used in conjunction with deficient substrate plasma, the APTT provides the basis for the quantitation of specific coagulation factors.

The intrinsic capability of blood to form a fibrin clot requires coagulation factors XII, XI, IX, VIII, X, V, II (prothrombin), fibrinogen, platelet lipid, and calcium.

By adding a substance to activate factors XII and XI, the contact factors, the partial thromboplastin time becomes the "activated" partial thromboplastin time (APTT). Because coagulation endpoints are shorter and sharper than with the PTT, the APTT has proven to be a simple and highly reliable measurement of the intrinsic coagulation mechanism.

5

TEST PROCEDURE

1. Preincubate 0.025M Calcium Chloride to 37°C.
2. Pipette 0.1 mL of reconstituted APTT reagent into a test cuvette.
3. Add 0.1 mL test or control plasma.
4. Incubate at 37°C for 5 minutes exactly.
5. Add 0.1 mL preincubated Calcium Chloride, simultaneously starting the timer.
6. Record the clotting time.

10

The prothrombin time is the method for monitoring oral anticoagulation therapy. It is also a fundamental screening test for a deficiency or abnormality of extrinsic coagulation factor VII, and the factors common to both the intrinsic and extrinsic hemostatic pathways: fibrinogen II, V and X. When used in conjunction with deficient substrate plasma, the PT provides the basis for the quantification of specific coagulation factors.

15

The capability of blood to form a fibrin clot by way of the extrinsic hemostatic pathway requires tissue thromboplastin, calcium, factor VII, factor V, factor X, factor II (prothrombin) and factor I (fibrinogen). When tissue thromboplastin and calcium are added to a sample of citrated plasma, the actions of the intrinsic factors are bypassed and the reaction becomes specific for the coagulation factors involved in the extrinsic and common pathways.

20

TEST PROCEDURE

1. Preincubate reconstituted PT reagent to 37°C.
2. Pipette 0.1 mL of test or control plasma into a test cuvette.
3. Incubate at 37°C for at least 2 minutes, but not more than 10 minutes.
4. Inject 0.2 mL of the preincubated reagent, simultaneously starting the timer.
5. Record the clotting time.

25

30

Template bleeding time measurements are obtained at the same time points as the blood samples, as mentioned previously. Template bleeding time is measured after a uniform incision is made on the mucous membrane of the inner upper lip with a Surgicutt[®] automated incision device (ITC, Edison, NJ). Blood is blotted with Surgicutt[®] bleeding time blotting paper every 30 seconds, being careful not to disturb the incision site. Template bleeding time is measured from the moment of the incision until the blood no longer stained the blotting paper. Bleeding times of 10 minutes are taken to be maximal.

35

Anti-Xa and anti-IIa activity

These samples are centrifuged at 1500 x g for 10 min at 4°C. The plasma is removed and stored at -70°C until assayed. Anti-Xa and Anti-IIa activity are analyzed by chromogenic methods using kits supplied by American Diagnostica (actichrome[®] Heparin and actichrome[®] Heparin anti-IIa, Greenwich, CT), with minor modifications. Incubations and reactions are performed at 37°C. Amidolytic activity (milli-optical units, or mOD) is determined using a SPECTRAMax microtiter plate spectrophotometer and Softmax Pro software (Molecular Devices Corp., Sunnyvale, CA). The 1st International LMWH Standard (National Institute for Biological Standards and Control, London; anti-Xa activity 168 IU/mg and anti-IIa activity 66.5 IU/mg) is used to construct standard curves for measuring heparin and ENOX anti-Xa and anti-IIa activity. The curves are constructed using a four parameter curve-fitting model (Softmax Pro, Molecular Devices, Sunnyvale, CA). Values for anti-Xa and anti-IIa activity of heparin and ENOX are reported in International Units.

Platelet aggregation

The Inhibition of ex-vivo Platelet Aggregation assay is based on that of Zucker, "Platelet Aggregation Measured by the Photoelectric Method", Methods in Enzymology 169, 117-133 (1989).

Platelet rich plasma (PRP) is prepared by centrifugation of the blood samples at 150 x g for 10 min. After removal of the supernatant containing PRP, platelet-poor plasma (PPP) is prepared by centrifugation of the remaining sample at 1000 x g for 10 min. Platelet count is determined with a Coulter ZM or Coulter ZBI particle counter (Coulter Instruments, Hialeah, FL). When necessary, platelet count is adjusted to 3×10^8 platelets/mL using autologous PPP. PRP (250 µL) is incubated at 37°C while being stirred at 1200 rpm. After preincubation with epinephrine for 1 min (1 µM, Chrono-par 393, Chrono-log Corp., Havertown, PA), platelet aggregation is induced by adenosine diphosphate (ADP, 10 µM, Chrono-par 384, Chrono-log Corp., Havertown, PA), collagen (equine tendon, 10 µg/mL, Chrono-par 385, Chrono-log Corp., Havertown, PA), arachidonic acid (1 mM, Biodata Corp, Horsham, PA) or thrombin (4 Units/mL, Enzyme Research Institute, South Bend, IN; plus Gly-Pro-Arg-Pro, a fibrin polymerization inhibitor, 2 mM, Sigma Chemical Co., St. Louis, MO). Platelet aggregation is monitored spectrophotometrically with a PAP-4C platelet aggregator (Bio Data Corp, Horsham, PA). Results are expressed as a percent inhibition of the rate of aggregation as compared to the pre-drug aggregation response.

Statistics

Data obtained by multiple sampling during the experiment are analyzed by two-way repeated measures analysis of variance. Post-hoc multiple comparisons of means to control values within treatment groups, and comparison of ENOX data to other treatment groups, are performed using the least significant difference test. A p-value less than 0.05 is considered significant.

Results

The combined use of an anti-Xa activity compound and a platelet aggregation antagonist compound used according to the invention, provides for use of those compounds at doses that would be subefficacious if used individually, while effecting the inhibition of repetitive platelet thrombus formation to the same extent as high doses of the platelet aggregation antagonist compound without significantly increasing template bleeding time (as compared to: a >5-fold increase in template bleeding time caused by high-dose of the platelet aggregation antagonist compound, BSPBPA; and a ~3-fold increase in template bleeding time caused by high-dose of the platelet aggregation antagonist compound, PBGACA).

Combination of a low-dose of a platelet aggregation antagonist compound with heparin, at a dose which increased APTT 2.0- to 2.5- fold over baseline, did not inhibit repetitive platelet thrombus formation.

The present invention may be embodied in other specific forms without departing from the spirit or essential attributes thereof.

WHAT IS CLAIMED IS:

1. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and pharmaceutically effective amounts of a compound having anti-Xa activity and a platelet aggregation antagonist compound.
5
2. The pharmaceutical composition according to claim 1 wherein the compound having anti-Xa activity is a low molecular weight heparin.
- 10 3. The pharmaceutical composition according to claim 1 wherein the low molecular weight heparin is selected from the group consisting of enoxaparin, nardroparin, dalteparin, certroparin, parnaparin, reviparin, ardeparin/RD heparin/ RDH or tinzaparin.
4. The pharmaceutical composition according to claim 1 wherein the low molecular weight
15 heparin is enoxaparin.
5. The pharmaceutical composition according to claim 1 wherein the compound having anti-Xa activity is a heparinoid compound.
- 20 6. The pharmaceutical composition according to claim 1 wherein the heparinoid compound is danaparoid.
7. The pharmaceutical composition according to claim 1 wherein the platelet aggregation antagonist compound is abciximab, N-[N-[N-(4-(piperidin-4-yl)butanoyl)-N-ethylglycyl]aspartyl]-
25 L- β -cyclohexyl alanine, N-[N-[N-(4-(piperidin-4-yl)butanoyl)-N-ethylglycyl]aspartyl]-L- β -cyclohexyl alanine amide or N-(n-butylsulfonyl)-4-(piperidin-4-yl)butyloxy-L-phenylalanine
8. The pharmaceutical composition according to claim 7 wherein the platelet aggregation antagonist compound antagonist is abciximab.
30
9. The pharmaceutical composition according to claim 7 wherein the platelet aggregation antagonist compound is N-[N-[N-(4-(piperidin-4-yl)butanoyl)-N-ethylglycyl]aspartyl]-L- β -cyclohexyl alanine.
- 35 10. The pharmaceutical composition according to claim 7 wherein the platelet aggregation antagonist compound is N-[N-[N-(4-(piperidin-4-yl)butanoyl)-N-ethylglycyl]aspartyl]-L- β -cyclohexyl alanine amide.

11. The pharmaceutical composition according to claim 7 wherein the platelet aggregation antagonist compound is N-(n-butylsulfonyl)-4-(piperidin-4-ylbutyloxy)-L-phenylalanine.

12. A method of treating or preventing a physiological condition associated with a thrombosis related ischemic disorder in a patient comprising administering to said patient pharmaceutically effective amounts of a compound having anti-Xa activity and a platelet aggregation antagonist compound.

13. The method of claim 12 wherein the physiological condition is selected from stroke, atherosclerosis, angiogenesis, thrombosis, a thromboembolic condition, disseminated intravascular coagulation, peripheral arterial occlusive disease, hemodialysis, an extra-corporeal circulation requirement associated with a surgical procedure, tissue damage caused by phospholipases A2, an acute coronary syndrome, or a thromboembolic syndrome associated with cancer, sepsis or obstetrical complications.

14. The method of claim 13 wherein the physiological condition is the acute coronary syndrome.

15. The method of claim 14 wherein the acute coronary syndrome is unstable angina or myocardial infarction.

16. The method of claim 12 wherein the treating or preventing occurs in the course of coronary artery bypass surgery or percutaneous transluminal coronary angioplasty.

17. The method of claim 12 wherein the treating or preventing occurs in the course of percutaneous transluminal coronary angioplasty.

18. The use of pharmaceutically effective amounts of a compound having anti-Xa activity and a platelet aggregation antagonist compound in the preparation of a medicament for treating or preventing a physiological condition associated with thrombosis related ischemic disorder.

19. The use according to claim 18, wherein the physiological condition is the acute coronary syndrome.

20. The use according to claim 19 wherein the acute coronary syndrome is unstable angina or myocardial infarction.

21. The use according to claim 18 wherein the treating or preventing occurs in the course of coronary artery bypass surgery or percutaneous transluminal coronary angioplasty.

22. The use according to claim 18 wherein the treating or preventing occurs in the course of percutaneous transluminal coronary angioplasty.

23. A kit for treating or preventing a physiological condition associated with thrombosis related ischemic disorder, said kit comprising a plurality of separate containers, wherein at least one of said containers contains a compound having anti-Xa activity and at least another of said containers contains a platelet aggregation antagonist compound, and said containers optionally contain a pharmaceutical carrier.

24. A kit according to claim 23 wherein of said containers at least one of said containers should contain the compound having anti-Xa activity without the presence of the platelet aggregation antagonist compound, and at least another of said containers should contain the platelet aggregation antagonist compound without the presence of the compound having anti-Xa activity.


25. A product containing a compound having anti-Xa activity and a platelet aggregation antagonist compound as a combined preparation for simultaneous, separate or sequential use for treating or preventing a physiological condition associated with thrombosis related ischemic disorder.

26. A use of a compound having anti-Xa activity and a platelet aggregation antagonist compound together in therapy for treating or preventing a physiological condition associated with thrombosis related ischemic disorder.

27. A pharmaceutical composition of a compound having anti-Xa activity and a platelet aggregation antagonist compound together in therapy for use in treating or preventing a physiological condition associated with thrombosis related ischemic disorder.

28. A pharmaceutical composition of a compound having anti-Xa activity and a platelet aggregation antagonist compound as described herein with respect to preferred embodiments.

FIG. 1



1 / 2 4

2 / 2 4

FIG. 2




FIG. 3
Effect of Coadministration of Enoxaparin or Heparin
with BSPBPA on APTT




FIG. 4

Effect of Coadministration of Enoxaparin or Heparin
with BSPBPA on PT





FIG. 5


Effect of Coadministration of Enoxaparin or Heparin with BSPBPA on Anti-Xa Activity



*: p<0.05 vs enoxaparin + BSPBPA

FIG. 6


Effect of Coadministration of Enoxaparin or Heparin with BSPBPA on Anti-IIa Activity



6 / 2 4

FIG. 7

Effect of Coadministration of Enoxaparin or Heparin with BSPBPA on Template Bleeding Time



7 / 2 4

FIG. 8

Effect of Coadministration of Enoxaparin or Heparin with BSPBPA on Platelet Count






FIG. 9

Effect of Coadministration of Enoxaparin or Heparin with BSPBPA on Collagen-induced *ex vivo* Platelet Aggregation



F I G. 10
Effect of Coadministration of Enoxaparin or Heparin with BSPBPA on ADP-induced ex vivo Platelet Aggregation



11 / 24

FIG. 11

Effect of Coadministration of Enoxaparin or Heparin with
BSPBPA on Arachidonic Acid-induced ex vivo Platelet




FIG. 12
Effect of Coadministration of Enoxaparin or Heparin with
BSPBPA on Thrombin-induced ex vivo Platelet Aggregation





FIG. 13



13 / 24

Blood Sampling/Hemodynamics

Arterial samples at -5, 5, 10, 30, 60 min


- APTT, PT
- Anti-Xa, Anti IIa
- Template Bleeding Time
- Platelet Count
- Platelet Aggregation (ADP, Collagen, Thrombin, AA)
- Coronary Blood Flow (CFRs per 20 min periods)

Treatment Groups (n=5 per group)

- PBGACA low (10 µg/kg + 0.15 µg/kg/min)
- PBGACA high (30 µg/kg + 0.45 µg/kg/min)
- Enoxaparin (0.5 mg/kg + 5 µg/kg/min)
- Enoxaparin (0.5 mg/kg + 5 µg/kg/min) plus PBGACA low
- Heparin (60 U/kg + 0.7 U/kg/min)
- Heparin (60 U/kg + 0.7 U/kg/min) plus PBGACA low

FIG. 14

Effect of Coadministration of Enoxaparin or Heparin
with PBGACA on CFRs



*: p<0.05 vs enoxaparin + PBGACA

14 / 24

FIG. 15

Effect of Coadministration of Enoxaparin or Heparin
with PBGACA on APTT




FIG. 16

Effect of Coadministration of Enoxaparin or Heparin
with PBGACA on PT

0' 5' 10' 30' 60'




FIG. 17

Effect of Coadministration of Enoxaparin or Heparin
with PBGACA on Anti-Xa Activity




FIG. 18
Effect of Coadministration of Enoxaparin or Heparin
with PBGACA on Anti-IIa Activity





FIG. 19

19 / 24

Effect of Coadministration of Enoxaparin or Heparin with PBGACA on Template Bleeding Time



*: p<0.05 vs enoxaparin + PBGACA

FIG. 20

Effect of Coadministration of Enoxaparin or Heparin
with PBGACA on Platelet Count




FIG. 2I

Effect of Coadministration of Enoxaparin or Heparin with
PBGACA on Collagen-induced ex vivo Platelet Aggregation




FIG. 22

Effect of Coadministration of Enoxaparin or Heparin with
PBGACA on ADP-induced ex vivo Platelet Aggregation





FIG. 23


Effect of Coadministration of Enoxaparin or Heparin with
PBGACA on Arachidonic Acid-induced ex vivo Platelet Aggregation



24 / 24

FIG. 24

Effect of Coadministration of Enoxaparin or Heparin with
PBGACA on Thrombin-induced ex vivo Platelet Aggregation



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/21440

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 31/725, 38/00, 31/445, 39/395, 39/40, 39/42

US CL :514/56, 18, 331; 424/143.1, 142.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/56, 18, 331; 424/143.1, 142.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Database Embase on STN, Elsevier Science B.V., AN 96261183, WENZL, M.E. et al. 'Loss of extremities in cases of heparin-induced thrombocytopenia,' abstract, Unfallchirurg (DE), 1996.	1-17, 23-25, 27
Y	US 5,292,756 A (DUGGAN et al.) 08 March 1994, the abstract and claims.	1-17, 23-25, 27
Y	Database Hcaplus on STN, American Chemical Society, AN 1995:886373 (Doc. No. 123:306599), WO 9510295 A1 (KLEIN, S.I. et al.), abstract.	1-17, 23-25, 27



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

15 MARCH 1998

Date of mailing of the international search report

06 APR 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

M. MORZIE

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/21440**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 18-22, 26
because they relate to subject matter not required to be searched by this Authority, namely:

Claims drawn to the "use of" a compound are non-statutory.
2. ☒ Claims Nos.: 28
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

Claim 28 is an omnibus-type claim, the metes and bounds of which are unclear.
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/21440

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

HCAPLUS, EMBASE-low molecular weight heparin compounds and heparinoid compounds herein (as anti xa compounds) and platelet aggregation antagonist compounds herein for the treatment of thrombotic disorders generally and particularly.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/21440

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Database Hcaplus on STN, American Chemical Society, AN 1996:236654 (Doc. No. 124: 331300), GENETTA, T.B., et al. 'Abciximab: A new antiaggregant used in angioplasty,' abstract, Ann. Pharmacother., 1996.	1-17, 23-25, 27
Y	Database Wpids on STN, Derwent Information Ltd., AN 96-049388 [05], WO 9534286 A1, (BENES, L.R.M. et al.), abstract.	1-17, 23-25, 27